

Pepper Cute Stuff

(*Capsicum annuum*)

Germination

Media

Use a well-drained, disease-free, soilless media with a pH of 5.8 to 6.2 and a medium initial nutrient charge (EC 0.75 mmhos/cm).

Sowing

Can be produced in 512 or larger plug tray. Cover the seed lightly with coarse vermiculite.

Stage 1 – Germination takes 5 to 7 days

Soil temperature: 71 to 78°F (22 to 24°C)

Light: Optional

Moisture: Keep soil wet (level 4) during Stage 1

Humidity: Maintain 95%+ relative humidity (RH) until radicles emerge

Note: Peppers are very sensitive to high salts, particularly high ammonium, during germination. Keep ammonium levels to less than 10 ppm.

Plug Production

Media

Use a well-drained, disease-free, soilless media with a pH of 5.8 to 6.2 and a medium initial nutrient charge (EC 0.75 mmhos/cm).

Sowing

Can be produced in 512 or larger plug tray. Cover the seed lightly with coarse vermiculite.

Stage 1 – Germination takes 5 to 7 days

Soil temperature: 71 to 78°F (22 to 24°C)

Light: Optional

Moisture: Keep soil wet (level 4) during Stage 1

Humidity: Maintain 95%+ relative humidity (RH) until radicles emerge

Note: Peppers are very sensitive to high salts, particularly high ammonium, during germination. Keep ammonium levels to less than 10 ppm.

Stage 2

Soil temperature: 68 to 73°F (20 to 23°C)

Light: Up to 2,500 f.c. (26,900 Lux)

Moisture: Reduce soil moisture slightly (level 3 to 4) to allow the soil to dry out slightly before watering for best germination and rooting.

Fertilizer: Apply fertilizer at rate 1 (less than 100 ppm N/less than 0.7 mS/cm EC) from nitrate-form fertilizers with low phosphorous.

Stage 3

Soil temperature: 65 to 70°F (18 to 21°C)

Light: Up to 2,500 f.c. (26,900 Lux)

Moisture: Allow media to dry further until the surface becomes light brown (level 2) before watering to promote root growth and control shoot growth. Keep the moisture level to wet-dry cycle (moisture level 4 to 2). Avoid permanent wilting.

Fertilizer: Increase fertilizer to rate 2 (100 to 175 ppm N/0.7 to 1.2 mS/cm EC) from 14-0-14 or other calcium/potassium nitrate fertilizer. Fertilize every 2 to 3 irrigations. Maintain medium pH of 5.8 to 6.2 and EC between 1.0 and 1.5 mS/cm (1:2 extraction).

Growth Regulators: If necessary, one application of Sumagic (uniconazole) at rate of 2.5ppm (4.6 ml/l of 0.055% formulation) can be applied at 2 weeks after sowing for Cajun Belle and Cute Stuff Red. Sweet Heat is a naturally compact variety and does not need Sumagic.

Stage 4

Soil temperature: 62 to 67°F (17 to 19°C)

Light: Up to 5,000 f.c. (53,800 Lux)

Moisture: Same as Stage 3.

Fertilizer: Fertilize with 14-0-14, 15-5-15 or calcium/potassium nitrate feed at rate 2 (100 to 175 ppm N/0.7 to 1.2 mS/cm EC) as needed.

Growing On to Finish

Container Size

4-in. (10-cm) square/quart pots: 1 plug per pot

Media

Use a well-drained, disease-free, soilless media with a pH of 5.5 to 6.2 and a medium initial nutrient charge (EC 0.75 mmhos/cm).

Temperature

Nights: 65 to 70°F (18 to 21°C)

Days: 68 to 79°F (20 to 26°C)

Peppers prefer warm temperatures. Peppers can be damaged by temperatures below 45°F (7°C).

Light

Maintain light levels as high as possible. High light levels and spacing will keep plants from stretching.

Irrigation

Grow plants on the dry side. To provide some height control, plants can be allowed to wilt slightly prior to irrigation once plants are rooted to the edge of the container.

Fertilizer

Apply fertilizer at rate 3 (175 to 225 ppm N/1.2 to 1.5 mS/cm) every third irrigation with predominately a nitrate-form fertilizer with low phosphorus and high potassium. Maintain media pH at 5.8 to 6.2.

For a constant fertilizer program, apply fertilizer at rate 2 (100 to 175 ppm N or 0.7 to 1.2 mS/cm).

Pinching

Pinching is not needed.

Crop Scheduling

Sow to transplant (512 cell plug): 5 to 6 weeks

Transplant to sale: 4 to 7 weeks

Common Problems

Insect: Watch for aphids

Disease: no severe disease

Garden and Landscape Information

- Plant in full sun after all danger of frost has passed.

Garden height, spread and fruit size:

Height: 21 in. (53 cm)

Spread: 20 in. (51 cm)

Fruit size: 3 in. (8 cm) x 2.5 in. (6 cm)

Note: Growers should use the information presented here as a starting point. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. Chemical and PGR recommendations are only guidelines. It is the responsibility of the applicator to read and follow all the current label directions for the specific chemical being used in accordance with all regulations.

PanAmericanSeed™