GrowerFacts

Cuphea Dynamite

(Cuphea ignea)

Plug crop time: 5 to 6 weeks Transplant to finish: 6 to 7 weeks Annuals Culture (revised 01/13/21)

- Well-suited to solo and mixed containers.
- · Also works in indoor plant programs.

General Information

| Exposure | Bloom Season | Height | Spread | Spacing |
|----------|--|------------------------|-------------------------|-------------------------|
| Sun | Spring, Late Spring, Summer, Late Summer | 8-10 in. (20-25 cm) | 10-12 in. (25-30 cm) | 10-14 in. (25-36 cm) |

Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media pH/EC (1:2) | Cover Seed |
|-----------|--------------------------|------------|-----------------|--------------------------------------|------------------------------|-------------|
| RAW | 288 | 3-4 | 5-6 | 4-6 | 5.5-5.8 pH 0.75 mmhos/cm | Light cover |

Plug Production

| | Stage 1 | Stage 2 | Stage 3 | Stage 4 |
|-------------|----------------------|---|-------------------------------------|---|
| Moisture | Level 4 | Level 3-4 | Level 3-4 | Level 3-4 |
| Temperature | 70-75°F (21-24°C) | | | |
| Light | Light | 2,500 f.c. (26,900 Lux) | 2,500 f.c. (26,900 Lux) | 2,500-5,000 f.c. (26,900-53,800 Lux) |
| Fertilizer | | Less than 100 ppm N (Less than 0.7 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) |

Propagation Key Tips

No PGRs required.

Growing on to Finish

| Growing on Temperature | Target Media pH/EC (1:2) | Fertilizer | Daylength |
|--|--------------------------------|------------|-------------|
| (day) 70-75°F (21-24°C) (night) 65-68°F (18-20°C) | 5.8-6.2 pH 1.2-1.5 mmhos/cm | | Day Neutral |

Crop Scheduling

| Container Size | Plugs/Pot | Crop Time | Season | PGR |
|---------------------|-----------|-------------|--------|-----|
| 4"/4.5"/Quart/10 cm | 1 (ppp) | 6-7 (weeks) | Spring | - |

Finishing Key Tips

No pinching required. Well-suited to solo and mixed containers. Also works in indoor plant programs.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures







Dynamite

nite Dynar