

Nemesia Poetry™ F₁ Series

(*Nemesia foetans*)

Annuals Culture (revised 01/12/21)

Plug crop time: 4 weeks

Transplant to finish: 5 to 7 weeks

- Uniform, grower-friendly varieties in bright colours are ideal for Spring and Autumn premium pack and 4-in./10-cm pot programs.
- Branches very well without pinching to produce bushy plants that fill out fast and full.
- Winter flowering will occur in milder temperatures; will also flower lightly in warmer conditions in partial shade.

NOTE: It is recommended to sow 4 pellets per cell.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Summer, Late Summer	12-14 in. (30-36 cm)	10-12 in. (25-30 cm)	6-8 in. (15-20 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PEL	288	4	4	4-5	5.5-6.2 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 2-3	Level 2-3
Temperature	65-70°F (18-21°C)	68-72°F (20-22°C)	68-72°F (20-22°C)	65-68°F (18-20°C)
Light	Optional	2,500 f.c. (26,900 Lux)	2,500 f.c. (26,900 Lux)	5,000 f.c. (53,800 Lux)
Fertilizer		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
PGR			daminozide/1,500-2,500 ppm/Spray	

Propagation Key Tips

Seedlings may be more stretchy if germinating under dark conditions. Do not use a growth regulator before radicle emergence, as this can delay or stop germination.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
(day) 62-68°F (17-20°C) (night) 55-62°F (13-17°C)	5.8-6.2 pH 0.7-1.0 mmhos/cm	100 to 175 ppm N (0.7 to 1.2 EC)	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	5-7 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	6-8 (weeks)	Spring	-
5"/6"/1 Gallon/15 cm	3 (ppp)	6-8 (weeks)	Spring	-

Common Problems

Insects: Check/monitor for Thrips, as they can spread INSV.

Finishing Key Tips

If necessary, use 2 to 3 applications of daminozide 2,500-5,000 ppm Spray.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures




Blue



Deep Pink



Pink



White



Mixture

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2022 Ball Horticultural Company